

REDUCING THE ENVIRONMENTAL IMPACT OF SOLVENTS FOR CATALYTIC REACTIONS

MARTYN POLIAKOFF and MICHAEL W. GEORGE

**School of Chemistry, University of Nottingham, University Park, Nottingham, NG7 2RD, United Kingdom*

The present state of research on catalysis for sustainable chemistry

Ever since the birth of Green Chemistry, there has been a focus on increasing the sustainability of the chemistry by reducing waste. Catalysis has since made impressive advances in maximizing the selectivity of reactions and minimizing the waste from stoichiometric reagents [1]. More recently, there have been further improvements by use of photo- and electrochemistry to carry out “reagentless reactions” [2]. Nevertheless, large volumes of waste in pharmaceutical, speciality and fine chemicals manufacture originate in the solvents used in the processes and for the purification of the products [3]. Not only can the solvents sometimes be toxic compounds, for example chlorinated organics, but most are hydrocarbon-based and pose flammability and environmental hazards. Even the use of water as a solvent can be problematic as it is costly to clean traces of organic substrates from the water before it can be safely discharged into the environment. Therefore, there remains a pressing need for reducing solvent waste to create more sustainable chemical manufacture via catalysis. Here we suggest one possible way forward, linked to efforts for mitigating climate change.

Our recent research contributions to catalysis for sustainable chemistry

We have pioneered the use of supercritical carbon dioxide (scCO₂) as a solvent for chemical reactions and devised methodologies for high pressure continuous catalytic reactions on laboratory and larger scales. CO₂ becomes supercritical above its critical point 30.9 °C and 73.8 bar. scCO₂ has relatively low solvent power (similar to that of an alkane) and, in general, solubility of a compound increases with pressure. However, scCO₂ has several potential advantages as a solvent: (i) CO₂ is very inexpensive compared to most solvents; (ii) solute separation merely requires release of the pressure and purification of the used CO₂ is simpler than purification of typical organic solvents; (iii) CO₂ is miscible with gaseous O₂ and is completely non-flammable, offering safety advantages for aerobic oxidation reactions [4]; (iv) CO₂ is also miscible with H₂ making it highly suitable as a solvent for hydrogenation. Therefore, our initial work focused on hydrogenation of relatively simple organic compounds with H₂ using a Pd heterogeneous

catalyst, with the reactions eventually being scaled up to a 1000-ton p.a. commercial plant [5], which was technically highly successful but was rendered uneconomic by steeply rising energy costs for recompressing the CO₂ (see below). However, we demonstrated that scCO₂ can offer additional advantages such as in the hydrogenation of levulinic acid to form γ -valerolactone where the high pressure phase behavior can be manipulated to separate pure product (Fig. 1) from the co-product H₂O with less energy than would be required for conventional separation by distillation [6].




Fig. 1. Schematic of the hydrogenation of levulinic acid (LA) over a ruthenium catalyst to form γ -valerolactone (GVL) in scCO₂ using H₂O as the co-solvent and exploiting the pressure-induced phase separation of GVL and H₂O to obtain pure GVL at the output of the reactor. The photos on the right show this phase separation, with a water-soluble dye, Direct Red 23, to highlight the separation of the water more clearly. For more details, see [6].

We have also combined some of the advantages of photochemistry [2] and scCO₂ to demonstrate that scCO₂ can be used for reactions with photo-generated singlet oxygen (¹O₂). The reactions of ¹O₂ are potentially attractive because of their high atom economy but, until now, there has been a reluctance to use them on a large scale because of the unstable nature of the intermediate products and the need to minimize the flammability of the solvent. Both of these problems can be addressed by use of scCO₂ in high-pressure flow photo-reactors, which have the additional advantage that ¹O₂ has a long lifetime in scCO₂ [7]. Although our reactors are still relative small, we have produced a variety of products using ¹O₂ in scCO₂ on scales that are at least an order of magnitude greater than have been previously reported [8]. One example, which involves both ¹O₂ and an unusual dual function heterogeneous catalyst, is the semi-synthesis of the anti-malarial drug artemisinin (Fig. 2) using toluene as a co-solvent and eliminating the need for toxic CF₃CO₂H, which is conventionally used as the acid catalyst in this reaction [9].

We have also shown how such scCO₂ flow reactors can easily be automated and combined with AI for self-optimizing catalytic reactions [10].




Fig. 2. The photo-oxidation of dihydroartemisinic acid, DHAA, by photo-generated singlet O₂ in scCO₂ with a toluene co-solvent to form Artemisinin with use of a bi-functional catalyst, the photosensitiser tetraphenylporphyrin, immobilized on the acid catalyst Amberlyst 15. Adapted from [9].

In addition, we have considered the wider environmental impact of making chemicals and, been inspired by the ideas of **Chemical Leasing** [11], which proposes that chemical manufacture should recognize the most customers buy chemicals for the effect that those chemicals produce (e.g. coating a surface, treating an illness, etc.). This does against the conventional business model that is based upon the assumption that you will earn more from selling more product. The supplier does not sell quantities Chemical Leasing but the supplier sells the performance i.e. the function of the chemical. Thus, by analogy with the electronics industry [12], we proposed Moore's Law for Chemistry, namely that *"sustainable chemists should strive to reduce the amount of a chemical needed to produce a given effect by a factor of two, for example within a 5 year cycle, and this process should be repeated for a number of cycles"* [13]. We build on this concept below.

Outlook: future developments of research on catalysis for sustainable chemistry

CO₂ is cheap, abundant and scCO₂ is a solvent with considerable potential for catalytic reactions. At the same time, scCO₂ is beset by the problem of high energy costs for compression of the CO₂. We believe that increasing concerns about climate change offer a real opportunity to resolve this problem. The opportunity lies at the core of some philosophical aspects of Green Chemistry, namely the need to expand the focus from design of chemical routes and selection of feedstocks to the wider problems of manufacturing, business models and supply chains [14]. Our proposal involves **Carbon**

Capture and Storage (CCS) which is increasingly recognized as a key technology, at least in the short term, for decarbonizing not only the energy sector but also other industries (e.g. cement making or steam reforming to make so-called ‘blue’ ammonia). However, a major barrier to implementation worldwide is that CCS introduces significant additional costs compared to releasing CO₂ to the atmosphere. Therefore, there is a real need for strategies which add value to the captured CO₂ by transforming it into a revenue-earning asset. Thus, one needs to address not only the environmental but also the economic pillar of the triple bottom line [15], a difference between sustainable and green chemistry.

The opening of the 1000-ton p.a. commercial plant [5] in 2002 mentioned above coincided with a large rise in energy prices, a problem which continues to reoccur and impact chemical and other manufacturing. This meant that the energy cost for compressing CO₂ rendered the plant uneconomic to operate. Since then, we have worked extensively in CCS and realize that the key is to combine CCS and scCO₂ solvents [16, 17]. That would make a huge difference because all of the compression costs will be met, as they are already an integral part of CCS. The novelty arises from linking significant potential revenue to what is currently a costly implementation of the net-zero agenda via CCS. It transforms captured CO₂ from a worthless waste to a valuable resource without the need to modify the CO₂ itself. This achieved by exploiting the compression energy stored in captured CO₂ thereby valorizing CO₂ by a physical process that is already inherent to the CCS technology. It tackles one of the key aspects within the circular economy of how to reduce solvent waste - one of the unsolved issues to drive sustainability noting the amount of CO₂ needed to be used for chemical industry is incredibly small compared to the vast amount that needs to be captured. Of course, the captured CO₂ is likely to contain impurities but we have demonstrated [17] that the predicted levels of impurities for CCS would be acceptable for the type of hydrogenation reaction that we have previously carried out on an industrial scale [5]. Most CCS systems are still at the development stage so that now is the right moment to take a systems view and combine CCS plants with sustainable chemical production by catalysis in scCO₂.

Moore’s Law for Chemistry (see above) [13] focuses on effect chemicals and, in many ways, catalysts are the ‘ultimate’ effect chemicals. They facilitate transformations that chemists would otherwise struggle to carry out efficiently. Therefore, we propose that, if catalysis is really to help ensure our future supply of chemicals in a sustainable way, we need to adopt and implement **Moore’s Law for Chemistry**. This will need both existing and new catalysts together with future developments to reduce not only the amount of catalysts required but also the quantity of chemicals needed to make those catalysts. That is we need to reduce over a 5-year cycle by 50% the chemical footprint of a process including the catalysts that we use in the future. This means not only reducing the amount of catalyst needed to promote a given transformation but also halving the amount of chemicals required to make those catalysts. And we need to repeat that 50% reduction

over several cycles so that catalysis as well as the whole process can sustainably fulfil the chemical needs of future generations.

We have presented two innovative ideas, namely (i) harnessing CCS to provide a source of CO₂ solvents which can be fed back into the CCS process after use for easy recycling and (ii) applying Moore's Law for Chemistry to catalysts and catalytic processes. Individually, each of these ideas could substantially increase the sustainability of chemical manufacture in the future. Linking them together could be transformative but there is still a need to apply to scCO₂ some of the innovations in reactors and processes that we, and others, have already made for continuous chemistry [2] so as to create scalable supercritical photochemical processes. Chemical and other manufacture is now in the throes of a 4th industrial revolution driven by digitization [18], which makes the application of Moore's Law to catalytic processes a realistically achievable goal for sustainable chemistry exploiting the ever increasing application of AI to manufacturing.

Acknowledgments

We thank the University of Nottingham and all of the agencies, charities and companies, who have funded our research. We are grateful to our collaborators, colleagues, co-workers, students and technicians for their contributions to our research.

References

1. R. A. Sheldon, *Green Chem.*, **19**, 18 (2017).
2. M. Poliakoff, M. W. George, *Phil. Trans. R. Soc. A*, **378**: 20190260 (2019).
3. P. J. Dunn, S. Galvin, K. Hettenbach, *Green Chem.*, **6**, 43 (2004).
4. G. R. Akien, A. O. Chapman, P. Licence, M. Poliakoff, *Green Chem.*, **12**, 310, (2010).
5. P. Licence, M. Sokolova, S. K. Ross, M. Poliakoff, *Green Chem.*, **5**, 99 (2003).
6. R. A. Bourne, J. G. Stevens, J. Ke, M. Poliakoff, *ChemComm* 4632 (2007).
7. R. A. Bourne, X. Han, M. Poliakoff, M. W. George, *Angew. Chem. Intl Ed.*, **48**, 5322 (2009).
8. L. Q. Wu, B. L. Abreu, A. J. Blake, L. J. Taylor, W. Lewis, *et al.*, *Org. Proc. Res. Dev.*, **25**, 1873 (2021).
9. Z. Amara, J. F. B. Bellamy, R. Horvath, S. J. Miller, *et al.*, *Nature Chem.*, **7**, 489 (2015).
10. R. A. Skilton, R. A. Bourne, Z. Amara, R. Horvath, J. Jin, *et al.*, *Nature Chem.* **7**, 1 (2015).
11. [Chemical Leasing | UNIDO](#) (accessed 19th Sept 2022)
12. D. C. Brock, "Understanding Moore's Law", Chem. Heritage Found., Philadelphia (2006).
13. M. Poliakoff, P. Licence, M. W. George, *Angew. Chem. Int. Ed.*, **57**, 12590 (2018).
14. [An Agenda to Mainstream Green Chemistry.pdf \(greenchemistryandcommerce.org\)](#) (2015) (accessed 19th Sept 2022).
15. A. Scerri, P. James, *Int. J. of Soc. Res. Methodology*, **13**, 41 (2010)
16. "Big Ideas for the Future" RCUK, London, (2011), page 45.
17. J. G. Stevens, P Gomez, R. A. Bourne, M. W. George *et al.*, *Green Chem.*, **13**, 2727, (2011).
18. P. Bellini, D. Cenni, N. Mitolo, P. Nesi, G. Pantaleo, *et al. J. Ind. Inf. Integration*, **26**, 100276 (2022).

Keywords for Indexing: Solvents, Supercritical, Carbon Dioxide, Hydrogenation, Oxidation, Singlet Oxygen, Photo-reactors, Carbon Capture & Storage, Chemical Leasing, Moore's Law for Chemistry.